

# TEST YOUR IRRIGATION WATER FOR PHYTOPHTHORA

*Phytophthora* is a fungus-like organism called water mold, that can infect a wide variety of nursery plant species, and it spreads in irrigation water.

*In this article, we will describe how you can test your irrigation water for Phytophthora.*



Surface water sources are often used for irrigating plants

## *Irrigation water can be a source of contamination*

Irrigation water demands vary from several hundred to several hundred thousand gallons of water per day, depending on the size of the nursery. Several sources of water may be used to meet these irrigation demands, including surface water (rivers, streams, canals, lakes, ponds, reservoirs), well water, rainwater, municipal drinking water, and recycled runoff water. Of these, only rainwater, well water, and municipal water are free of pathogens. You should assume that all other sources are contaminated and disinfest them before use.

Alternatively, you can test your water sources at frequent intervals and treat only if necessary. Once pathogens gain entry into the nursery, they can spread through the irrigation system, infect plants, accumulate in the runoff water, and establish in the water storage reservoirs. Disinfecting contaminated water before it is used for irrigation is essential for breaking this cycle.

### *Web resources*

1. **Phytophthora host list:** <https://pnwhandbooks.org>
2. **Baiting for Phytophthora video:** <http://cleanwater3.org/decontaminate.asp#decontaminatetab2>
3. **Pocket Diagnostics® Phytophthora rapid kit:**
  - a. **Outside USA:** [www.pocketdiagnostic.com/onlineshop/](http://www.pocketdiagnostic.com/onlineshop/)
  - b. **Within USA:** [www.uidaho.edu/cals/parma-research-and-extension-center/plant-pathology](http://www.uidaho.edu/cals/parma-research-and-extension-center/plant-pathology)

## *Phytophthora*

There are more than 100 *Phytophthora* species; some are host-specific, while others can cause disease on hundreds of plant species. All require water to complete their life cycle.

*Phytophthora* species cause foliar blight, stem canker, shoot dieback and root rot on ornamentals, native plants, forest trees and agricultural crops worldwide. Many nursery plant species are highly susceptible, such as azalea, rhododendron, boxwood, and many other conifer species<sup>1</sup>.

The plant nursery environment is optimal for growth and proliferation of *Phytophthora*. Oospores and chlamydospores are capable of long-term survival in soil or infected plant debris. Under wet conditions, *Phytophthora* sporangia release numerous small swimming spores called zoospores, which can infect plants and cause disease. The zoospores not only survive and propagate in surface water but also can be delivered throughout the nursery in irrigation water.




*Phytophthora* sporangium releasing zoospores (top left, photo courtesy of Fred Schwenk), foliar blight symptom on rhododendron plants (top right), and rhododendron root rot caused by *Phytophthora* (bottom).

## *Baiting can be used to test irrigation water for Phytophthora*

### *Baiting technique*

Baiting is a relatively simple technique that uses susceptible plant parts as “baits” to attract *Phytophthora* spp. This method selects for live, active zoospores, which can swim toward and infect the bait.

You can use rhododendron leaves or hard (unripe) green pears (Bartlett, d' Anjou) as bait to detect *Phytophthora*, as they are susceptible to many species of *Phytophthora*. These baits can also capture several *Pythium* species, which are closely related to *Phytophthora* and cause damping-off of seedlings and stem cutting rots. There are two basic ways to bait: in a resealable bag (indoors), or in the water body itself (outdoors).



Baiting of water (top) with rhododendron leaves (left) and pear (right) baits, and the resulting brown lesions (bottom) caused by *Phytophthora* species.

### *Baiting indoors*

Collect a quart of water in a clean, soap-free container from near the surface of the water you wish to test. Pour half the water into each of two 1-gallon resealable bags labeled with the date and sample name. Place a rhododendron leaf or a hard green pear in each bag. Be sure to choose leaves or fruits without spots, blemishes or bruises; they should be free from pesticides that could interfere with the test. It is always a good idea to place a few leaves in bags with tap water or distilled water as a negative control. Allow the bags to sit for seven days at 65 to 75°F out of direct sunlight. Then remove the leaves or the pears and examine them for chocolate-brown spots (“lesions”) on rhododendron leaves or brownish red circular lesions on pears. If the pear has been sitting upright, you can sometimes see a “bathtub ring” of lesions around the pear made by zoospores. Spots made by *Phytophthora* are firm to the touch; if they are soft and mushy, they are likely caused by a soft rot organism; not *Phytophthora*. For more information, watch the tutorial on baiting for *Phytophthora* at the CleanWater3 website<sup>2</sup>.

### *Baiting outdoors*

You will need to make a mesh bag to hold the leaf baits. Plastic window screen material works well for this purpose. Some mesh bags are made with sleeves for individual leaves, as shown. Attach the mesh bag to an air-filled, plastic milk jug to act as a float, as the greatest number of zoospores will be near the water surface. Secure the mesh bag with a nylon rope, so you can cast the bag out into the water and then retrieve it seven days later. Remove the leaves from the bag and examine them for dark, chocolate-brown lesions. The outdoor baiting works well when the water temperature is between 48°F to 71°F.

If *Phytophthora* is present in water, zoospores will swim toward the surface of water surrounding the bait, colonize it and develop dark brown lesions. If the tested water is contaminated with a high amount of the pathogen, then brown lesions may appear within two to three days after baits are suspended in water. The appearance of lesions on the bait is a good indication that *Phytophthora* species are present, but this should be confirmed.

Rhododendron leaves in a mesh bag showing lesions after baiting outdoors (photo courtesy of Marianne Elliott)



---

## Detecting *Phytophthora* after baiting

To confirm the presence of *Phytophthora*, you can remove a small piece of the bait lesion and test it with a commercially available diagnostic kit for *Phytophthora* species. *Phytophthora* diagnostic kits are based on polyclonal antibodies for detection of multiple *Phytophthora* species. The kits are rapid, easy to use and relatively inexpensive. For example, the Pocket Diagnostics® *Phytophthora* kit<sup>3</sup> costs about \$8 each when purchased in boxes of 50 kits, and it takes less than 10 minutes to conduct the test. These kits are designed for genus-level detection of *Phytophthora* species, but will not identify the particular species.



Lateral flow device showing positive (left) and negative (right) results for *Phytophthora* detection using the Pocket Diagnostics *Phytophthora* rapid test kit.

If the *Phytophthora* test is positive and you would like to know what species is present, you can send the bait for additional testing by a university plant diagnostic lab (see next page). Note that it is much easier to send leaf baits rather than pears through the mail! Be sure to contact the lab in advance to make sure they can handle your sample. Plant diagnostic labs can attempt to identify the species by growing it in a petri dish containing an agar nutrient medium selective for *Phytophthora*. Identification can be based on microscopic features or DNA-based techniques. For most growers, genus-level identification is enough to demonstrate that their water is contaminated and requires treatment.

## Once you know if your water is infested, you can take steps to disinfect it.

Irrigation water treatment can minimize the spread of *Phytophthora* and help reduce losses. There are several methods for water disinfection with chemical, physical and biological modes of action. You can choose which method to use based on the mode of action, the volume of water to be treated, installation and operational costs involved, space requirements, and safety and environmental concerns. You can refer to the *waterborne solutions tool* that summarizes water treatment technologies used to control plant pathogens at the CleanWater3 website: <http://cleanwater3.org/growertools.asp>.

## A success story in managing irrigation water to reduce *Phytophthora*

A large container nursery in Oregon recycles 90 percent of its irrigation water; it succeeds in preventing *Phytophthora* infestation of its irrigation water by treating it first with sodium hypochlorite or calcium hypochlorite. We collected water samples from several different steps along the irrigation water pathway, and baited the water samples with rhododendron leaves to see if *Phytophthora* was present. We then used a DNA sequencing technique to identify the *Phytophthora* species that were on the bait leaves. The diagram (Fig. 1) illustrates results of baiting irrigation water in the month of September 2015.

---

## Plant disease diagnostic labs offering services to test for *Phytophthora*

### **Plant Diagnostics Laboratory at University of Massachusetts**

[www.ag.umass.edu/services/plant-diagnostics-laboratory](http://www.ag.umass.edu/services/plant-diagnostics-laboratory)

Baiting water for *Pythium*, *Phytophthora*, and *Rhizoctonia* **\$50**

### **Florida Extension Plant Diagnostic Clinics at University of Florida**

[www.plantpath.ifas.ufl.edu/extension/plant-diagnostic-center](http://www.plantpath.ifas.ufl.edu/extension/plant-diagnostic-center)

Baiting for *Pythium* or *Phytophthora* **\$65**

### **The Pullman Plant Pest Diagnostic Clinic at Washington State University**

[www.plantpath.wsu.edu/diagnostics](http://www.plantpath.wsu.edu/diagnostics)

Baiting water for *Phytophthora* **\$40**

### **Oregon State University Plant Clinic**

[www.plant-clinic.bpp.oregonstate.edu](http://www.plant-clinic.bpp.oregonstate.edu)

Baiting water for *Phytophthora* or *Pythium* **\$75**

### **Plant Disease Diagnostic Clinic at Cornell University**

[www.plantclinic.cornell.edu](http://www.plantclinic.cornell.edu)

- ELISA assay for *Phytophthora* **\$70**
- Immunostrip test for *Phytophthora* **\$60**
- PCR for *Phytophthora ramorum* **\$50**

### **Plant Diagnostic Lab at North Dakota State University**

[www.ag.ndsu.edu/pdl/services-and-fees](http://www.ag.ndsu.edu/pdl/services-and-fees)

Immunostrip test for *Phytophthora* **\$45**

### **Plant Disease Diagnostic Clinic at University of Wisconsin-Madison**

[www.pddc.wisc.edu/services-fees](http://www.pddc.wisc.edu/services-fees)

Testing for *Aphanomyces*, *Pythium* or *Phytophthora* **\$35**

### **Texas Plant Disease Diagnostic Lab at Texas A & M University**

[www.plantclinic.tamu.edu/fees](http://www.plantclinic.tamu.edu/fees)


ELISA assay for *Phytophthora* **\$20**

*continued....*

The main source of water for the nursery was a creek. The creek was infested with four *Phytophthora* spp., including two pathogenic species complexes, *P. parsiana* and *P. megasperma*. (A complex is a group of closely related species that cannot be distinguished with our method of sequencing.) The creek water was pumped to a retention reservoir where we detected two species complexes: the *P. parsiana* complex from the creek, and the *P. citricola* complex from another source, likely the runoff water. From there, the water was filtered and chlorinated, with a target concentration of 2 ppm and a 10-minute contact time. No *Phytophthora* species were detected in the

chlorinated water, which was then used for overhead irrigation in the growing areas of the nursery. Although the irrigation water was clean, and the plants did not show any symptoms of *Phytophthora*, we did detect *P. citricola* complex in the soil/crushed rock material underneath the containers.

Soil in nurseries is commonly infested by *Phytophthora* species, which can survive for years in bits of plant debris that infiltrate the soil/crushed rock. It is very difficult to disinfect contaminated soil. Runoff water from these growing areas, after contacting the contaminated soil, was found to harbor *P. parsiana* complex, *P. citrophthora*, and the *P. citricola* complex. Pumping the runoff water back into the retention reservoir carried these *Phytophthora* species with it. Fortunately, the subsequent chlorination treatment prevented *Phytophthora* from entering the irrigation water, so the contamination cycle was broken.



Chlorination reduces *Phytophthora* propagules circulating in irrigation water at a container nursery in Oregon.

Following this example, you can test your irrigation water along its flow path at regular (monthly) time intervals. By routinely testing the water with baiting, you can assess your risk for waterborne *Phytophthora*, and can implement preventive measures, if necessary.

Copyright © 2018 Neelam R. Redekar and Jennifer L. Parke  
All rights reserved.

**Acknowledgments:** Funding for this material is based upon work that is supported by the National Institute of Food and Agriculture, U. S. Department of Agriculture, Specialty Crop Research Initiative competitive program under award number 2014-51181-22372.

**Disclaimer:** Inclusion of a commercial product in the article reported here implies no endorsement by the authors or Oregon State University.

**Dr. Neelam R. Redekar** is a Postdoctoral Scholar in the Department of Crop and Soil Science at Oregon State University, Corvallis, Oregon; she can be reached at [Neelam.Redekar@oregonstate.edu](mailto:Neelam.Redekar@oregonstate.edu)

**Dr. Jennifer L. Parke** is a Professor (Senior Research) in the Department of Crop and Soil Science and the Department of Botany and Plant Pathology at Oregon State University in Corvallis, Oregon; she can be reached at [Jennifer.Parke@oregonstate.edu](mailto:Jennifer.Parke@oregonstate.edu)